

Google

puppels fiber wavenumber

Search

Advanced Search
Preferences

Web

Results 1 - 10 of about 412 for **puppels fiber wavenumber**. (0.35 seconds)

Discriminating basal cell carcinoma from perilesional skin using ...

In contrast, the use of the high wave-number region allows for **fiber optic probes ...** and G.J. **Puppels**, "Tissue characterization using high wavenumber Raman ...
link.aip.org/link/?JBOPFO/12/034004/1 - Similar pages

Tissue characterization using high wave number Raman spectroscopy ...

L. F. Santos, R. Wolthuis, S. Koljenović, R. M. Almeida, and G. J. **Puppels**, "High wavenumber fiber-optic probes for in vivo Raman spectroscopy" ...
link.aip.org/link/?JBOPFO/10/031116/1 - Similar pages

Glasses and Thin Films for Nanophotonics and Integrated Optics

... S. Koljenovic, Rui M. Almeida and Gerwin J. **Puppels**, "Fiber-optic Probes for in Vivo Raman Spectroscopy in the High-Wavenumber Region", Anal. Chem. ...
www.demat.ist.utl.pt/glasses/photronics - 26k - Cached - Similar pages

Fiber-optic probes for in vivo raman spectroscopy in the high ...

Fiber-optic probes for in vivo raman spectroscopy in the high-wavenumber region ...
KOLJENOVIC S. (2); ALMEIDA Rui M. (1); **PUPPELS** Gerwin J. (2); ...
cat.inist.fr/?aModele=afficheN&cpsidt=17233597 - Similar pages

Fiber-Optic Probes for in Vivo Raman Spectroscopy in the High ...

Fiber-Optic Probes for In Vivo Raman Spectroscopy in the High-Wavenumber Region ... S. Koljenovi , Rui M. Almeida, and Gerwin J. **Puppels** ...
pubs.acs.org/cgi-bin/abstract.cgi/ancham/2005/77/i20/abs/ac0505730.html - Similar pages

Puppels GJ

Puppels GJ (Gerwin J) In contrast, the use of the high wave-number region allows for fiber optic probes with a very simple design. ...
lib.biolinfo.pl/auth:**Puppels GJ** - 50k - Cached - Similar pages

Raman spectroscopy and its urological applications Hanchanale VS ...

Santos LF, Wolthuis R, Koljenović S, Almeida RM, **Puppels GJ**. Fiber-optic probes for in vivo Raman spectroscopy in the high-wavenumber region. ...
www.indianjrol.com/preprintarticle.asp?id=39550 - 73k - Cached - Similar pages

Tissue characterization using high wave number Raman spectroscopy.

Fused silica has no Raman signal in the high wave number region (2400-3800 cm⁻¹). This enables the use of a single unfiltered fiber to guide laser light ...
www.ncbi.nlm.nih.gov/pubmed/16229641 - Similar pages

Biomedical applications of Raman and infrared spectroscopy to ...

[28] L.F. Santos, R. Wolthuis, S. Koljenovic, R.M. Almeida and G.J. **Puppels**, Fiber-optic probes for in vivo Raman spec-. troscopy in the high-wavenumber ...
iospress.metapress.com/index/2UY07C4PD2K6NE8T.pdf - Similar pages

Methodology for fiber-optic Raman mapping and FTIR imaging of ...

detected successfully by analysis of the high-wavenumber. region. This suggests that development of an unfiltered. single fiber-optic probe for Raman ...
www.springerlink.com/index/N07648109H362637.pdf - Similar pages

puppels fiber wavenumber

Search

[Search within results](#) | [Language Tools](#) | [Search Tips](#) | [Dissatisfied? Help us improve](#) | [Try Google Experimental](#)
